

The statistics of diffusive flux

Alba Margarita Reséndiz Antonio⁽¹⁾, Hernán Larralde⁽²⁾

(1) *Facultad de Ciencias, UAEMor. Av Universidad s/n,*

CP 62210 Cuernavaca Morelos, Mexico

(2) *Centro de Ciencias Físicas, UNAM; Apartado postal 48-3,*

CP 62251 Cuernavaca Morelos, Mexico

We calculate the explicit probability distribution function for the flux between sites a in a simple discrete-time diffusive system composed by independent random walkers. We highlight some of the features of the distribution and we discuss its relation to the local instantaneous entropy production in the system. Our results are applicable both to equilibrium and non equilibrium steady states as well as for certain time dependent situations.

PACS numbers:

Random walkers are used extensively as phenomenological models for diffusive processes[1]. The connection between physical diffusion and random walks arises from the fact that the concentration in a system containing many identical independent random walkers satisfies Fick's law. That is, the mean flux between sites is proportional to the difference in the mean number of walkers at each site, plus, perhaps, a convective term if the walks are not symmetric. However, to the best of our knowledge, the statistics of the flux in such a system have not been characterized (beyond the mean, of course). This is not the case in more complicated systems, for which a large amount of work on the statistics of currents has been carried out. Recent examples include statistics and large deviation theory for the fluctuations of the current in lattice gases [2, 3, 4]; the integrated current distribution for the steady-state of the one-dimensional zero-range process [5]; the joint probability function for the occupation number and the current through the system in the asymmetric simple exclusion process with open boundaries [6], to mention but a few.

In addition to being an alternative approach to the description of transport properties of simple diffusive systems, the statistics of flux are of relevance in many diffusion limited processes. For example, in a diffusion reaction system in which the reactants are initially separated, fluctuations in the flux of reactants into the interfacial region give rise to fluctu-

ations in the position of the reaction front and in the net reaction kinetics [7, 8, 9].

Furthermore, viewed from the thermodynamic side, there has been a great deal of work relating the statistics of flux to the entropy production of non equilibrium steady states [11, 12, 13]. From this perspective, diffusive systems modeled by random walkers are among the simplest examples that can be used to test and extend such ideas.

In this work we obtain an explicit expression for the probability distribution of the flux, that is, the net number of particles that hop between neighboring sites in a given time step, in a system containing independent discrete time random walkers. This system has the added bonus that the results are applicable to both equilibrium and non-equilibrium situations, where the latter can be achieved by imposing concentration differences on the boundaries or by subjecting the system to an external field (by considering biased random walkers) or both.

For definiteness, in this work we consider a discrete one dimensional system in which independent discrete time random walkers evolve synchronously according to the following rules: at each time step, a walker moves to the site on its right or to the site on its left with probabilities p and q respectively, or stays at its site with probability $r = 1 - p - q$. We note that the discreteness in time allows us to consider fluxes that arise from the simultaneous hopping of many particles. Such multiparticle events are not present in continuous time versions of the system; we discuss the continuous limit of the problem further on.

The system in which the random walkers evolve consists of $l + 2$ sites and we impose as boundary conditions that the number of walkers at sites $i = 0$ and $i = l + 1$ be Poisson distributed with fixed mean values c_0 and c_{l+1} respectively [10].

To compute the probability distribution of particle flux between neighboring sites, labeled i and $i + 1$, we require J_+ and J_- , the number of particles that jump from site i to site $i + 1$ and the number of particles that jump from $i + 1$ to i respectively. In terms of these, the total flux J between these sites will be given by $J = J_+ - J_-$.

We denote by m and n the occupancy of sites i and $i + 1$ respectively. Then, the probability that the net flux between these sites is J can be expressed as

$$p_i(J) = \sum_{m,n} p(J|m,n)p_{i,i+1}(m,n), \quad (1)$$

where $p_{i,i+1}(m,n)$ is the probability of finding exactly m and n particles at sites i and $i + 1$, and $p(J|m,n)$ is the probability of having a total flux J between these sites given those occupancy numbers.

Since the walkers are independent, we have

$$p(J|m, n) = \sum_{J_-=0}^{\infty} p_+(J + J_-|m) p_-(J_-|n), \quad (2)$$

where

$$p_+(J_+|m) = \binom{m}{J_+} p^{J_+} (1-p)^{m-J_+} \quad (3)$$

and

$$p_-(J_-|n) = \binom{n}{J_-} q^{J_-} (1-q)^{n-J_-} \quad (4)$$

are the probabilities that J_+ particles jump to the right from a site containing m particles and J_- particles jump to the left from a site containing n particles.

In what follows, it will prove useful to work with the generating function of expression (2), defined as $\hat{p}(z|m, n) := \sum_{J=-\infty}^{\infty} z^J p(J|m, n)$, which is given by

$$\hat{p}(z|m, n) = \left(1 - q + \frac{q}{z}\right)^n (1 - p + zp)^m. \quad (5)$$

Next, we note that this system has the remarkable property that the joint probability distribution $p_{i,i+1}(m, n)$ of the occupancy numbers factorizes into a product of Poisson distributions [15]. This property is analogous to the factorization of the steady state occupancy distribution which occurs in certain zero range processes under similar boundary conditions [16]. Thus, the joint occupancy distribution $p_{i,i+1}(m, n)$ can be written as

$$p_{i,i+1}(m, n) = \varphi_i(m) \varphi_{i+1}(n), \quad (6)$$

where

$$\varphi_i(M) = \frac{1}{M!} e^{-c_i} c_i^M, \quad (7)$$

and c_i is the mean number of particles at site i . Actually, under certain circumstances, namely initial conditions already characterized by independent Poisson distributions, the joint occupancy distribution can be expressed as the product of Poisson distributions throughout the evolution of the system. In this case, the mean occupation numbers can be obtained as the solution to the discrete diffusion equation

$$c_i(t+1) - c_i(t) = p c_{i-1}(t) + q c_{i+1}(t) - (p+q) c_i(t) \quad (8)$$

with appropriate initial and boundary conditions. However, just as for zero range processes [16], the system always reaches a unique factorizable steady state, independently of the initial conditions, for which the parameters c_i are given by

$$c_i = \frac{1}{\left(\frac{p}{q}\right)^{l+1} - 1} \left[c_0 \left(\frac{p}{q}\right)^{l+1} - c_{l+1} + (c_{l+1} - c_0) \left(\frac{p}{q}\right)^i \right], \quad (9)$$

which is the steady state solution of equation (8).

Thus, we are in a position to evaluate the generating function $\hat{p}_i(z)$ of the distribution of fluxes between sites i and $i+1$:

$$\hat{p}_i(z) = \sum_m \sum_n \hat{p}(z|m, n) \varphi_i(m) \varphi_{i+1}(n) = e^{-pc_i - qc_{i+1} + zpc_i + \frac{q}{z}c_{i+1}}. \quad (10)$$

From this expression, the moments and the cumulants of the distribution may be readily evaluated. For example, all the odd cumulants are given by

$$\kappa_{2n-1} = pc_i - qc_{i+1}, \quad n = 1, 2, \dots \quad (11)$$

whereas the even cumulants are

$$\kappa_{2n} = pc_i + qc_{i+1}, \quad n = 1, 2, \dots \quad (12)$$

In particular, using the explicit expression for the concentration profile for the steady states given in equation (9), we find

$$\kappa_{2n-1} = \frac{1}{p^{l+1} - q^{l+1}} [(p - q)(c_0 p^{l+1} - c_{l+1} q^{l+1})] \quad (13)$$

and

$$\kappa_{2n} = \frac{1}{p^{l+1} - q^{l+1}} [(p + q)(c_0 p^{l+1} - c_{l+1} q^{l+1}) + 2pq(c_{l+1} - c_0)p^i q^{l-i}] \quad (14)$$

Furthermore, using the explicit expressions for the moments we can calculate the skewness γ_1 and excess (or kurtosis) γ_2 of the distribution, which are given by

$$\gamma_1 = \frac{\langle (J - \langle J \rangle)^3 \rangle}{\sigma^3} = \frac{(pc_i - qc_{i+1})}{(pc_i + qc_{i+1})^{\frac{3}{2}}} \quad (15)$$

and

$$\gamma_2 = \frac{\langle (J - \langle J \rangle)^4 \rangle}{\sigma^4} - 3 = \frac{1}{(pc_i + qc_{i+1})}. \quad (16)$$

Thus, only in the limit $pc_i + qc_{i+1} \gg 1$ is the distribution of flux essentially Gaussian. In fact, the expression for $\hat{p}_i(z)$ can be inverted to obtain an explicit expression for $p_i(J)$. Rewriting

$$\hat{p}_i(z) = e^{-pc_i - qc_{i+1}} \exp \left[\frac{1}{2} (2\sqrt{pqc_i c_{i+1}}) \left(z \sqrt{\frac{pc_i}{qc_{i+1}}} + \frac{1}{z \sqrt{\frac{pc_i}{qc_{i+1}}}} \right) \right], \quad (17)$$

and comparing with the generating function of the modified Bessel functions [14]

$$\exp \left[\frac{1}{2} x \left(t + \frac{1}{t} \right) \right] = \sum_{s=-\infty}^{\infty} I_s(x) t^s, \quad (18)$$

yields

$$p_i(J) = e^{-pc_i - qc_{i+1}} \left[\frac{pc_i}{qc_{i+1}} \right]^{\frac{J}{2}} I_J(2\sqrt{pqc_i c_{i+1}}). \quad (19)$$

This distribution, shown in Fig.1, shares some of the properties of the Gaussian distribution: It is characterized by only two parameters, say pc_i and qc_{i+1} in this case. It satisfies a generalized stability condition, in the sense that the distribution of the sum of independent (integer) random variables distributed according to $p_i(j)$, has the same form as $p_i(j)$ with appropriately rescaled parameters. That this is the case is obvious from the explicit expression of the generating function, but it could have been foreseen from the actual process we are describing. Indeed, the complete derivation presented above applies also for the distribution of flux between adjacent cell in systems of arbitrary dimension. For these, the one dimensional system we are considering corresponds to a projection onto one of the axes. Thus, the flux distribution we obtain can itself be thought of as that of the sum of many independent fluxes with distributions similar to $p_i(j)$.

The large $|J|$ behavior of $p_i(J)$, which is where it most clearly differs from the Gaussian, is easily amenable to evaluation. From the series expansion of the Bessel functions [14], we immediately find that

$$p_i(J) \sim \begin{cases} e^{-pc_i - qc_{i+1}} \frac{(pc_i)^J}{J!} \left(1 + \frac{pqc_i c_{i+1}}{J} + \dots \right) & \text{when } J \rightarrow \infty \\ e^{-pc_i - qc_{i+1}} \frac{(qc_{i+1})^{|J|}}{|J|!} \left(1 + \frac{pqc_i c_{i+1}}{|J|} + \dots \right) & \text{when } J \rightarrow -\infty. \end{cases} \quad (20)$$

Another limiting behavior of $p_i(J)$ worth mentioning is that corresponding to the continuous time limit. This limit is achieved by assuming that the hopping probabilities can be expressed as $p = \alpha \delta t$ and $q = \beta \delta t$, where δt is the time interval between successive steps.




FIG. 1: Semilogarithmic plots of $p_i(J)$ with variance equal to 2 and a) $\langle J \rangle = 1$; b) $\langle J \rangle = 0$ (dot-dashed lines). Gaussian distributions with the same variance and mean as $p_i(j)$ in each case are plotted for comparison.

Then, keeping only the terms up to linear order in δt we find

$$p_i(J) \approx \begin{cases} \alpha c_i \delta t & J = 1 \\ 1 - (\alpha c_i + \beta c_{i+1}) \delta t & J = 0 \\ \beta c_{i+1} \delta t & J = -1. \end{cases} \quad (21)$$

All other values of J have probabilities of higher order in δt and are, therefore, negligible as $\delta t \rightarrow 0$. The above expression merely reflects the fact that in the continuum limit, the transport of particles amongst neighboring sites during a time interval δt is, at most, a single particle process. The statistical parameters in this limit can be calculated directly from the expressions obtained above; thus, for example, the mean number of particles that flow from site i to site $i + 1$ in a time interval δt is $(\alpha c_i - \beta c_{i+1}) \delta t$ as was to be expected.

A further characteristic that $p_i(J)$ shares with the Gaussian distribution is that the ratio $p_i(J)/p_i(-J)$ is a pure exponential. This quantity is of interest because it can be associated to entropy production of non-equilibrium steady states. Indeed, given the explicit form of $p_i(J)$ between sites i and $i + 1$ in a system, and the symmetry properties of the Bessel functions, we have:

$$s_i = \ln \left(\frac{p_i(J)}{p_i(-J)} \right) = J \ln \left(\frac{p c_i}{q c_{i+1}} \right) \quad (22)$$

The average of this quantity is

$$\langle s_i \rangle = (pc_i - qc_{i+1}) \ln \left(\frac{pc_i}{qc_{i+1}} \right) \quad (23)$$

which has the familiar non-negative form that appears in the H -theorem, and that we argue can be considered as an instantaneous local entropy production for this system. It should be remarked that this is not the same quantity that appears in the fluctuation theorems [11, 12, 13], which are concerned with the large fluctuations of the time-integrated flux.

Also, from the definition of s_i in equation (22) it is again apparent that this quantity is a random variable, the statistics of which are simply related to the statistics of flux described above. Thus, for example, the variance of s_i will be given by

$$\langle s_i^2 \rangle - \langle s_i \rangle^2 = (pc_i + qc_{i+1}) \left[\ln \left(\frac{pc_i}{qc_{i+1}} \right) \right]^2 \quad (24)$$

To justify our identification of $\langle s_i \rangle$ as an average local entropy production, first note that the equilibrium for this system is defined as the steady state with zero average flux. In this situation, the concentration profile will be given by

$$c_i = c_0 (p/q)^i, \quad (25)$$

where c_0 is the concentration imposed on the boundary site $i = 0$. On the other hand, from the thermodynamic perspective the system corresponds to noninteracting particles in an external field h . For such a system the concentration profile is given by the barometric equation

$$c_i = e^{\beta(\mu - ih)}, \quad (26)$$

where μ is the chemical potential and β is the inverse temperature. Comparing both expressions for the concentration profiles at equilibrium leads to the identification

$$\beta\mu = \ln c_0; \quad \beta h = \ln \frac{p}{q}. \quad (27)$$

Of course, in general the system is not in equilibrium; however, since the occupancy is characterized by independent Poisson distributions, it can be considered as being in local equilibrium. Thus, we can rewrite equation (23) as:

$$\langle s_i \rangle = \langle J \rangle_i \left[\ln c_i - \ln c_{i+1} + \ln \frac{p}{q} \right] = -\langle J \rangle_i [\Delta_i \beta (\mu_i + \Phi(i))], \quad (28)$$

where $\Phi(i) = -ih$ is a “potential energy” and Δ_i is the difference operator. Equation (28) is a discrete analog of the local entropy production of linear thermodynamics.

At this point it is worth emphasizing that the average entropy production proposed in equation (23) is obtained from a *single* step local statistic. Thus, this quantity should be interpreted as the entropy produced at that time step, averaged over an ensemble of statistically identical systems, which is the usual interpretation of statistical mechanics. This is especially relevant in the time dependent case, for which all of the above results also hold, as mentioned above, given that the initially the sites of each system of the ensemble are occupied according to independent Poisson distributions. For these it makes no sense to study time averaged quantities of single systems and relate the results to the ensemble averages. On the contrary, in the steady state the ensemble average of s_i , Eq. (22), will coincide with the time average of the flux in a single system, yielding the connection usually assumed by the ergodic hypothesis. This will not be true for higher moments of s_i due to correlations in the flux at different times.

If we restrict ourselves to the steady states, the average flux is constant and we can calculate the total entropy production in the system as

$$\begin{aligned} s_{\text{total}} &= \sum_{i=0}^l \langle s \rangle_i = \sum_{i=0}^l (pc_i - qc_{i+1}) \ln \left(\frac{pc_i}{qc_{i+1}} \right) \\ &= \frac{1}{p^{l+1} - q^{l+1}} [(p - q)(c_0 p^{l+1} - c_{l+1} q^{l+1})] [(l + 1) \ln(p/q) + \ln(c_0/c_{l+1})], \end{aligned} \quad (29)$$

where c_0 and c_{l+1} are the concentrations imposed on the boundaries of the system and we used the explicit expression for the steady state flux given in equation (9). In particular, if we take the limit $q = p$, the above expression coincides with that obtained in [15], where the entropy production is interpreted in terms of a time asymmetry in the dynamical randomness between the forward and backward paths of the diffusion process [15, 17].

Finally, the limit in which $q \rightarrow 0$ say, is worth discussing. This totally asymmetric hopping case is again of some importance in the context of zero range processes. The distribution of flux in this case is easy to evaluate from the explicit expression in equation (19), from which we obtain $p_i(J) = 0$ for $J < 0$ and $p_i(J)$ is Poissonian with mean pc_i for $J > 0$. Further, as discussed above, a field can be associated to the $\ln(p/q)$, which gives rise to a “potential energy”; thus, in the limit when q vanishes, motion against the field is impossible and motion along the field “dissipates” an infinite amount of energy. In accordance, the mean entropy production diverges in this limit.

In summary, we have obtained the explicit probability distribution function of single step flux between adjacent sites in a diffusive system composed of independent discrete random walkers. The key to solving the problem is the fact that the joint site occupancy distribution

for this system factorizes into independent Poisson distributions even in non equilibrium situations. We also obtain the statistical parameters that characterize the distribution and discuss some of its limiting behaviors. In addition, we use the distribution to evaluate a single step local entropy production, whose average can be related to the usual expressions from nonequilibrium thermodynamics. In this context, an interesting extension of this work would be to introduce an additional conserved quantity which can be distributed among the random walkers at each site. This energy-like quantity may or may not affect the jumping rates (although from a physical perspective it probably should), and would allow the study of coupled transport phenomena in these extremely simple settings. We have thus far been unable to pose such scenario in a tractable manner. A different avenue of research would be to determine multiple time and site flux statistics in systems containing many random walkers, characterizing, for example, the correlations in flux at two sites of the system as a function of time.

This work was partially supported by grant IN-100803 of DGAPA UNAM. We thank D. Sanders, F. Leyvraz and M. Aldana for the many useful comments and suggestions on this manuscript.

APPENDIX

The key result that permits the explicit calculation of $p_i(J)$ is that the joint occupation probability distribution of the system factorizes into Poisson distributions. For the sake of completeness, we show how this comes about.

We require the determination of the joint occupation probability distribution, that is, the probability $p_t(n_0, n_1, \dots, n_{l+1})$ of finding $n_0, n_1, \dots, n_{l+1}$ particles, at the sites $0, 1, \dots, l+1$ respectively at time t , with the boundary conditions described in the text. The evolution equation for this quantity is

$$\begin{aligned}
 p_{t+1}(n_0, n_1, \dots, n_{l+1}) &= e^{-c_0} \frac{c_0^{n_0}}{n_0!} e^{-c_{l+1}} \frac{c_{l+1}^{n_{l+1}}}{n_{l+1}!} \int p_t(m_0, m_1, \dots, m_{l+1}) \\
 &\times \prod_{i=1}^l \delta(m + l_{i-1}^+ + l_{i+1}^- - (l_i^+ + l_i^-) - n_i) \\
 &\times \prod_{i=0}^{l+1} \binom{m_i}{l_i^-} \binom{m_i - l_i^-}{l_i^+} q^{l_i^-} p^{l_i^+} r^{m_i - l_i^- - l_i^+},
 \end{aligned} \tag{30}$$

where

$$\int = \sum_{m_0=0}^{\infty} \dots \sum_{m_l=0}^{\infty} \sum_{m_{l+1}=0}^{\infty} \sum_{l_0^-=0}^{\infty} \dots \sum_{l_l^-=0}^{\infty} \sum_{l_{l+1}^-=0}^{\infty} \sum_{l_0^+=0}^{\infty} \sum_{l_1^+=0}^{\infty} \dots \sum_{l_{l+1}^+=0}^{\infty}$$

and we recall that $r = 1 - p - q$. Equation (31) looks messy, but it is actually very easy to understand: if we consider the LHS of this equation without the delta functions, we note that it contains all the possible events from all the possible configurations at time t , weighted by the probability that the events occur. The deltas restrict this immense sum to the terms which end up in the occupancy configuration specified on the RHS of the equation.

In expression (31) we have already imposed Poisson occupancy distributions with concentrations c_0 and c_{l+1} on the sites 0 and $l+1$ as boundary conditions for the system. This will enable us to drive the system into “nonequilibrium” steady states by imposing different concentrations on the boundary, or considering biased random walks (i.e. $p \neq q$), or both.

This evolution equation can be dealt with by evaluating the multivariate generating function of the joint distribution, namely:

$$\hat{p}_t(z_0, \dots, z_{l+1}) = \sum_{n_0=0}^{\infty} \sum_{n_1=0}^{\infty} \dots \sum_{n_{l+1}=0}^{\infty} z_0^{n_0} z_1^{n_1} \dots z_{l+1}^{n_{l+1}} p_t(n_0, n_1, \dots, n_{l+1}). \quad (31)$$

After some simple but tedious algebra, which consists in regrouping terms and adding up binomial expansions, the generating function can be shown to satisfy the recursion relation

$$\begin{aligned} \hat{p}_{t+1}(z_0, \dots, z_{l+1}) &= e^{-c_0(1-z_0)} e^{-c_{l+1}(1-z_{l+1})} \\ &\times \sum_{m_0=0}^{\infty} \dots \sum_{m_{l+1}=0}^{\infty} p_t(m_0, m_1, \dots, m_{l+1}) \\ &\times [1 + (z_1 - 1)p]^{m_0} [q + z_2p + z_1r]^{m_1} \\ &\times [z_{l-1}q + p + z_lr]^{m_l} [1 + (z_l - 1)q]^{m_{l+1}} \\ &\times \prod_{i=2}^{l-1} [z_{i+1}p + z_{i-1}q + z_ir]^{m_i}. \end{aligned} \quad (32)$$

Comparing with the definition of the generating function, we note that this expression is equivalent to

$$\begin{aligned} \hat{p}_{t+1}(z_0, z_1, \dots, z_l, z_{l+1}) &= e^{-c_0(1-z_0)} e^{-c_{l+1}(1-z_{l+1})} \times \\ &\hat{p}_t(1 + (z_1 - 1)p, \quad z_2p + q + z_1r, \quad z_3p + z_1q + z_2r, \dots \\ &\dots, \quad z_lr + z_{l-2}q + z_{l-1}r, \quad z_{l-1}q + p + z_lr, \quad 1 + (z_l - 1)q). \end{aligned} \quad (33)$$

To solve this equation we make the ansatz that the distribution is given by the product of Poisson distributions, and thus, its generating function can be written as

$$\hat{p}_t(z_0, \dots, z_{l+1}) = e^{-\sum_{i=0}^{l+1} c_i(t)(1-z_i)}. \quad (34)$$

Substitution of this expression into Eq.(33) leads to the following equations for the mean occupation numbers $c_i(t)$ which characterize the Poisson distributions at each site:

$$\begin{aligned} \sum_{i=0}^{l+1} c_i(t+1)[1-z_i] &= c_0[1-z_0] + c_{l+1}[1-z_{l+1}] + c_0(t)p[1-z_1] + c_1(t)[1-(z_2p+q+z_1r)] \\ &+ \sum_{i=2}^{l-1} c_i(t)[1-(z_{i+1}p+z_{i-1}q+z_ir)] + c_l(t)[1-(z_{l-1}q+p+z_lr)] + c_{l+1}(t)q[1-z_l] \end{aligned}$$

If we now impose the boundary conditions $c_0(t) = c_0$ and $c_{l+1}(t) = c_{l+1}$ for all times t , the above equation may be rewritten as

$$\begin{aligned} \sum_{i=1}^l c_i(t+1) - \sum_{i=1}^l c_i(t) &= (c_0p - c_1(t)q) - (c_l(t)p - c_{l+1}q) \\ &+ \sum_{i=1}^l [c_i(t+1) - c_{i-1}(t)p - c_{i+1}(t)q - c_i(t) + c_i(t)(p+q)]z_i \end{aligned} \quad (35)$$

which must be valid for all values of the variables z_i . This is achieved if, in addition to the boundary conditions, the parameters $c_i(t)$ satisfy the discrete diffusion equation

$$c_i(t+1) - c_i(t) = pc_{i-1}(t) + qc_{i+1}(t) - (p+q)c_i(t). \quad (36)$$

Thus, if we have an ensemble of systems in which each site is initially occupied according to independent Poisson distributions, then the occupancy distribution will remain being the product of Poisson distributions throughout the evolution of the process.

Furthermore, independently of the initial conditions, the system will reach a unique stationary state characterized by the product distribution with the parameters c_i solutions of the stationary state of equation (36):

$$c_i = \frac{1}{\left(\frac{p}{q}\right)^{l+1} - 1} \left[c_0 \left(\frac{p}{q}\right)^{l+1} - c_{l+1} + (c_{l+1} - c_0) \left(\frac{p}{q}\right)^i \right]. \quad (37)$$

With this expression, we can calculate the explicit values of the mean flux and its variance as functions of p , q , the imposed boundary concentrations c_0 and c_{l+1} , and, for the variance, the position along the system:

$$\langle J_N \rangle = \frac{1}{p^{l+1} - q^{l+1}} [(p-q)(c_0p^{l+1} - c_{l+1}q^{l+1})] \quad (38)$$

$$\sigma_i^2 = \frac{(p+q)}{(p-q)} \langle J_N \rangle + 2pq \left(\frac{c_{l+1} - c_0}{p^{l+1} - q^{l+1}} \right) p^i q^{l-i}. \quad (39)$$

Finally, it is worth mentioning that the above derivation can also be extended to include site dependent jump rates.

-
- [1] G. H. Weiss *Aspects and Applications of the Random Walk* North-Holland, (1994)
 - [2] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim; Phys. Rev. Lett. 94, 030601 (2005)
 - [3] T. Bodineau and B. Derrida Phys. Rev. Lett. 92, 180601 (2004)
 - [4] B. Derrida, B. Douot, and P.-E. Roche; Journal of Statistical Physics, Vol. 115, Nos. 3/4, May 2004
 - [5] R. J. Harris, A Rákos and G M Schütz arXiv:cond-mat/0504711
 - [6] M. Depken and R. Stinchcombe Phys. Rev. Lett. 93 040602-1 2004
 - [7] S. M. Cox and M. D. Finn Phys. Rev. E 63, 051102 (2001)
 - [8] M. Sinder and J. Pelleg Phys. Rev. E 62, 3340-3348 (2000)
 - [9] M. Araujo, H. Larralde, S. Havlin, and H. E. Stanley, Phys. Rev. Lett. 71, 3592 (1993)
 - [10] This is a natural choice of boundary conditions if we wish to model contacts with a chemical baths at equilibrium, for which the number of particles is Poisson distributed in the grand canonical ensemble. Further, as shown in the appendix, this choice allows the exact factorizability of the joint occupancy distribution.
 - [11] G. Gallavotti and E. G. D. Cohen, Phys. Rev. Lett. 74, 2694 (1995).
 - [12] J. Kurchan, J. Phys. A: Math. Gen. 31, 3719 (1998).
 - [13] J. L. Lebowitz and H. Spohn, J. Stat. Phys. 95, 333 (1999)
 - [14] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, Dover (1972)
 - [15] P. Gaspard New Journal of Physics 7 (2005) 77
 - [16] E. Levine, D. Mukamel, and G.M. Schütz; arXiv:cond-mat/0412129
 - [17] P. Gaspard Journal of Statistical Physics, Vol. 117, Nos. 3/4, November 2004